

Exercise session 6

1. Let $\mathcal{F} = \{x \in \mathbb{R}^n \mid g_i(x) \leq 0, i = 1, \dots, m\}$ where g_i are convex functions on \mathbb{R}^n . Show that \mathcal{F} is a convex set.

2. Let f be a convex function on some convex set C . Let $x_1, \dots, x_n \in C$ and $\lambda_1, \dots, \lambda_n$ such that $\lambda_i \geq 0$ for all $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n \lambda_i = 1$. Show Jensen's inequality

$$f\left(\sum_{i=1}^n \lambda_i x_i\right) \leq \sum_{i=1}^n \lambda_i f(x_i)$$

for $n = 3$.

3. Prove the arithmetic-geometric mean inequality for positive x_1, \dots, x_n , i.e.

$$\frac{x_1 + \dots + x_n}{n} \geq (x_1 \cdot \dots \cdot x_n)^{1/n}.$$

Use Jensen's inequality and the convex function $-\log$.

4. Is $h(x) = |x| + \max\{e^x, 10 + 37x + x^6\}$ a convex function on \mathbb{R} ?

5. a) Show that $g(x) = x^3$ is not a convex function on \mathbb{R} .

b) Find a convex domain $C \subset \mathbb{R}$ such that $g(x) = x^3$ is a convex function on C .

7. Problem 3 from the SF1811 exam 13-01-2016.