

A Characterization of $(3 + 1)$ -Free Posets

Mark Skandera

*Department of Mathematics, Massachusetts Institute of Technology, Cambridge,
 Massachusetts 02139*

E-mail: skan@math.mit.edu

Communicated by the Managing Editors

Received January 25, 1999

Posets containing no subposet isomorphic to the disjoint sums of chains $3 + 1$ and/or $2 + 2$ are known to have many special properties. However, while posets free of $2 + 2$ and posets free of both $2 + 2$ and $3 + 1$ may be characterized as interval orders, no such characterization is known for posets free of only $3 + 1$. We give here a characterization of $(3 + 1)$ -free posets in terms of their antiadjacency matrices. Using results about totally positive matrices, we show that this characterization leads to a simple proof that the chain polynomial of a $(3 + 1)$ -free poset has only real zeros. © 2001 Academic Press

Key Words: poset; f -vector; total positivity.

1. INTRODUCTION

For nonnegative integers a and b , we denote by $\mathbf{a} + \mathbf{b}$ the poset which is the disjoint sum of an a -element chain and a b -element chain. A poset is called $(\mathbf{a} + \mathbf{b})$ -free if it contains no induced subposet isomorphic to $\mathbf{a} + \mathbf{b}$. (See [13, Chap. 3] for basic definitions.) For example, the first two posets in Fig. 1 are $2 + 2$ and $3 + 1$. The third poset P is $(2 + 2)$ -free but not $(3 + 1)$ -free, because the subposet induced by the elements $\{2, 3, 4, 6\}$ is isomorphic to $3 + 1$. For any poset P , we will denote the order relation in P by the symbol $<_P$, reserving $<$ for comparisons of numbers and vectors.

Fishburn [6] characterized $(2 + 2)$ -free posets by showing that any such poset P may be represented as a set of closed intervals of real numbers $[c_i, d_i]$, ordered by

$$[c_i, d_i] <_P [c_j, d_j], \quad \text{if } d_i < c_j.$$

He also showed that for posets free of both $3 + 1$ and $2 + 2$, the intervals may be chosen to have unit length. In honor of these results, posets in the above classes are often called *interval orders* and *unit interval orders*.




FIGURE 1

A second well known characterization of $(2+2)$ -free, $(3+1)$ -free posets involves natural labellings and antiadjacency matrices. Let P be a poset with n elements. Any bijective function $\phi: P \rightarrow \{1, \dots, n\}$ is called a *labelling* of P , and is called *natural* if it satisfies $\phi(x) < \phi(y)$ whenever $x <_P y$. If the elements x and y are labelled as i and j , we will often write $i <_P j$ to mean $x <_P y$. For a fixed labelling of P , we define an *antiadjacency matrix* $A = [a_{ij}]$ by

$$a_{ij} = \begin{cases} 0, & \text{if } i <_P j, \\ 1 & \text{otherwise.} \end{cases}$$

THEOREM 1.1. *Let P be a poset on n elements. P is $(2+2)$ -free and $(3+1)$ -free if and only if it may be naturally labelled so that the corresponding antiadjacency matrix A satisfies $a_{jk} \geq a_{i\ell}$, for all integers $1 \leq i \leq j \leq n$ and $1 \leq k \leq \ell \leq n$.*

That is, the positions (i, j) such that $a_{ij} = 0$ form a Ferrers shape in the upper right corner of A .

Unfortunately, little is known about posets free *only* of $3+1$. Characterization of $(3+1)$ -free posets is desirable, because several results and conjectures about posets require avoidance only of $3+1$. For instance, Stanley's generalization of the chromatic polynomial [14] is known to be s -positive for the incomparability graphs of $(3+1)$ -free posets [10], and is conjectured to be e -positive for these graphs as well [14, 17, 19]. Further, the chain polynomial of a $(3+1)$ -free poset has only real zeros. (See [17] and Corollary 4.1.) This implies that the f -vector of a $(3+1)$ -free poset is log-concave and unimodal.

We will characterize $(3+1)$ -free posets with a result analogous to Theorem 1.1. Observing several properties of squared antiadjacency

matrices in Section 2, we use these properties to prove the main theorem in Section 3. In Section 4 we derive a result of Stanley [17, Corollary 2.9] as a corollary of the main theorem and discuss some open questions regarding the chain polynomials of posets.

2. THE SQUARED ANTIADJACENCY MATRIX

Let P be a finite poset with elements labelled $\{1, \dots, n\}$, and let A be the corresponding antiadjacency matrix. The *squared antiadjacency matrix* $B = A^2$ has a simple combinatorial interpretation. Let $G = (P, E)$ be the graph whose *adjacency matrix* is A . The vertex set of G consists of the elements $\{1, \dots, n\}$ of P , and the edge set consists of the ordered pairs $(i, j) \in P \times P$ such that $i \not\leq_P j$. (See Fig. 2.) Clearly, $B = [b_{ij}]$ counts paths of length two in G . That is, b_{ij} is the number of ordered triples (i, x, j) , where (i, x) and (x, j) belong to E .

OBSERVATION 2.1. Assume that P is $(3+1)$ -free, and let i, j, k , and ℓ be elements of P .

- (1) If $i <_P j <_P k$, then $b_{ik} = 0$.
- (2) If $b_{ik} > b_{i\ell}$, then there is an element $x <_P \ell$, such that $i \not\leq_P x \not\leq_P k$.
- (3) If $b_{ik} > b_{jk}$, then there is an element $y >_P j$, such that $k \not\leq_P y \not\leq_P i$.

Proof. (1) Assume $b_{ik} > 0$. Then, for some element x of P , (i, x) and (x, k) belong to E , implying that $i \not\leq_P x \not\leq_P k$. In fact, x must be incomparable to i and k , for if $x <_P i$, then $x <_P i <_P j <_P k$, and if $k <_P x$, then




FIGURE 2

$i <_P j <_P k <_P x$, both impossible. Similarly, x cannot be comparable to j . Thus, the subposet of P induced by $\{i, j, k, x\}$ is isomorphic to $\mathbf{3} + \mathbf{1}$, contradicting our assumption that P is $(\mathbf{3} + \mathbf{1})$ -free.

(2) If $b_{ik} > b_{i\ell}$, then there are more paths of length two in G from i to k than from i to ℓ . It follows that for some element x of P , the pairs (i, x) and (x, k) belong to E , and the pair (x, ℓ) does not.

(3) Apply the argument of (2) to the dual poset of P . ■

Elements such as x in Observation 2.1(2) are central to the proof of Lemma 2.2. To simplify notation, we introduce the following definition.

DEFINITION 2.1. Let i, k, ℓ , and x be elements of P . Call x a (k, ℓ) -*advantage* for i if (i, x) and (x, k) are edges in the graph G and (x, ℓ) is not.

We use the word *advantage*, imagining that x helps us to travel from i to k , but not from i to ℓ . In the language of partially ordered sets, x is a (k, ℓ) -advantage for i if $x <_P \ell$, and $i \not\leq_P x \not\leq_P k$. Note that in Fig. 2, the vertex x is a (k, ℓ) -advantage for i , although it is not a (k, ℓ) -advantage for j .

LEMMA 2.2. Assume that P is $(\mathbf{3} + \mathbf{1})$ -free, and let i, j, k , and ℓ be elements of P . If $b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}$, then one of the following is true:

- (1) P contains an element x such that $j <_P x <_P \ell$ and $b_{j\ell} = 0$.
- (2) P contains an element y such that $i <_P y <_P k$ and $b_{ik} = 0$.

Proof. Let us denote by $\alpha(k, \ell, i)$ the number of elements of P which are (k, ℓ) -advantages for i . Then,

$$b_{ik} - b_{i\ell} = \alpha(k, \ell, i) - \alpha(\ell, k, i).$$

Assuming that $b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}$, we have

$$\alpha(k, \ell, i) + \alpha(\ell, k, j) > \alpha(k, \ell, j) + \alpha(\ell, k, i),$$

and at least one of the following two inequalities must be true,

$$\alpha(k, \ell, i) > \alpha(k, \ell, j),$$

$$\alpha(\ell, k, j) > \alpha(\ell, k, i).$$

Suppose that $\alpha(k, \ell, i) > \alpha(k, \ell, j)$. Then P contains an element x which is a (k, ℓ) -advantage for i and not a (k, ℓ) -advantage for j . By Definition 2.1, the pairs (i, x) and (x, k) belong to E and the pairs (x, ℓ) and (j, x) do not. Thus, $j <_P x <_P \ell$ and by Observation 2.1(1), $b_{j\ell} = 0$. Similarly, if $\alpha(\ell, k, j) > \alpha(\ell, k, i)$, then P contains an element y such that the pairs (y, k) and (i, y) do not belong to E . Thus, $i <_P y <_P k$ and $b_{ik} = 0$.

3. MAIN RESULT

THEOREM 3.1. *Let P be a poset on n elements. P is $(\mathbf{3} + \mathbf{1})$ -free if and only if it may be naturally labelled so that the squared antiadjacency matrix B satisfies the following two conditions for all integers $1 \leq i \leq j \leq n$ and $1 \leq k \leq \ell \leq n$.*

- (1) $b_{jk} \geq b_{i\ell}$
- (2) If $b_{ik} - b_{i\ell} \neq b_{jk} - b_{j\ell}$, then $b_{i\ell} = 0$ and $b_{ik} < b_{jk} - b_{j\ell}$.

EXAMPLE 3.1. Corresponding to any natural labelling of the poset P in Fig. 2 is the squared antiadjacency matrix

$$B = \begin{bmatrix} 2 & 2 & 0 & 0 & 0 \\ 2 & 2 & 0 & 0 & 0 \\ 4 & 4 & 2 & 2 & 0 \\ 4 & 4 & 2 & 2 & 0 \\ 5 & 5 & 3 & 3 & 1 \end{bmatrix}.$$

Condition (1) of the theorem says that entries of the squared antiadjacency matrix increase to the left in rows and downward in columns. To prove that a $(\mathbf{3} + \mathbf{1})$ -free poset P may be labelled to satisfy this condition, we will begin with an arbitrary labelling of P and the squared antiadjacency matrix B . Note that relabelling P corresponds to simultaneous row and column permutation of B . Let us denote the i th row and i th column of any matrix M by M_{i*} and M_{*i} .

OBSERVATION 3.2. *The following two conditions on any real matrix M are equivalent.*

(1) *It is possible to simultaneously permute the columns and rows of M so that its entries weakly increase to the left in rows and downward in columns.*

(2) *The rows and columns of M corresponding to any pair of indices i and j satisfy one of the following pairs of vector inequalities.*

- (a) $M_{i*} \geq M_{j*}$ and $M_{*i} \leq M_{*j}$.
- (b) $M_{i*} \leq M_{j*}$ and $M_{*i} \geq M_{*j}$.

The first statement simply says that we may sort the columns of M in weakly decreasing order while simultaneously sorting the rows in weakly increasing order. With a moment's thought, we see that this is possible if and only if the conditions in the second statement are true.

PROPOSITION 3.3. *Any $(\mathbf{3} + \mathbf{1})$ -free poset may be naturally labelled so that the entries of its squared antiadjacency matrix weakly increase to the left in rows and downward in columns.*

Proof. Let P be a $(\mathbf{3} + \mathbf{1})$ -free poset and assume that for each labelling of P , the corresponding antiadjacency matrix fails to satisfy condition (2) of Observation 3.2. (Trivially, condition (2) fails to hold for each non-natural labelling of P .) We consider two cases for a fixed labelling of P and the corresponding squared antiadjacency matrix B .

Case 1. Two columns of B are incomparable as vectors. That is,

$$B_{*i} \not\geq B_{*j} \quad \text{and} \quad B_{*i} \not\leq B_{*j},$$

for some indices $i \neq j$. Then for some indices $k \neq \ell$ we have $b_{ik} > b_{i\ell}$ and $b_{jk} < b_{j\ell}$, implying that

$$b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}.$$

Applying Lemma 2.2 to this inequality, we have $b_{j\ell} = 0$ or $b_{ik} = 0$, both contradictions. The argument for incomparable rows is identical.

Case 2. All columns of B are pairwise comparable as vectors, as are all rows, but for some indices $i \neq j$, we have an incorrect pair of comparisons of the form

$$B_{i*} \not\geq B_{j*} \quad \text{and} \quad B_{*i} \not\geq B_{*j}.$$

That is, there are elements k and ℓ in P , not necessarily distinct, satisfying $b_{i\ell} > b_{j\ell}$ and $b_{ki} > b_{kj}$.

By Observation 2.1(2), P contains an element $x <_P j$, such that $k \not\prec_P x \prec_P i$. By Observation 2.1(3), P contains an element $y >_P j$, such that $i \not\prec_P y \prec_P \ell$. Thus, $x <_P j <_P y$ is a chain, and each of these three elements is incomparable to i . This contradicts our assumption that P is $(\mathbf{3} + \mathbf{1})$ -free. ■

It is not hard to show that any labelling of a $(\mathbf{3} + \mathbf{1})$ -free poset which satisfies the first condition of Theorem 3.1 also satisfies the second condition.

PROPOSITION 3.4. *Let P be a $(\mathbf{3} + \mathbf{1})$ -free poset, naturally labelled so that its squared antiadjacency matrix B weakly increases to the left in rows and downward in columns. Let i, j, k , and ℓ be numbers satisfying $1 \leq i < j \leq n$ and $1 \leq k < \ell \leq n$. Then the 2×2 submatrix*

$$\begin{bmatrix} b_{ik} & b_{i\ell} \\ b_{jk} & b_{j\ell} \end{bmatrix}$$

satisfies one of the following two conditions:

- (1) $b_{ik} - b_{i\ell} = b_{jk} - b_{j\ell}$.
- (2) $b_{i\ell} = 0$ and $b_{ik} < b_{jk} - b_{j\ell}$.

Proof. Suppose that condition (1) is not satisfied.

Case 1. $(b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell})$. We apply Lemma 2.2 to this inequality. If $b_{j\ell} = 0$, then $b_{i\ell} = 0$ and $b_{ik} > b_{jk}$, contradicting our assumptions about weakly increasing entries of B . If instead $b_{ik} = 0$, then $b_{i\ell} = 0$ and $b_{j\ell} > b_{jk}$, another contradiction.

Case 2. $(b_{ik} - b_{i\ell} < b_{jk} - b_{j\ell})$. Again we apply Lemma 2.2. If $b_{jk} = 0$, then all four numbers are zero, a contradiction. We conclude that $b_{i\ell} = 0$ and that condition (2) is satisfied. ■

Finally, we show that the only posets satisfying the conditions of Theorem 3.1 are those which are $(3+1)$ -free.

PROPOSITION 3.5. *Let P be a labelled poset containing $3+1$ as an induced subposet, and let B be its squared antiadjacency matrix. Then there are two distinct elements i and k such that*

$$b_{ik} \neq 0 \quad \text{and} \quad b_{ii} - b_{ik} \neq b_{ki} - b_{kk}.$$

Proof. Let 1, 2, 3, and 4 be four elements of P such that $1 <_P 2 <_P 3$ is a chain, and 4 is incomparable to 1, 2, and 3. Let $G = (P, E)$ be the graph defined in Section 2.

Clearly, $b_{13} \neq 0$, since $(1, 4)$ and $(4, 3)$ are edges in G . We claim that

$$b_{11} - b_{13} \neq b_{31} - b_{33}.$$

Define the sets

$$X = \{x \in P \mid (1, x) \in E, (x, 1) \in E, (x, 3) \notin E\},$$

$$Y = \{x \in P \mid (x, 1) \in E, (x, 3) \notin E\},$$

and note that

$$|X| = b_{11} - b_{13},$$

$$|Y| = b_{31} - b_{33}.$$

Certainly X is a subset of Y . Moreover, it is a proper subset, since the element 2 belongs to Y and not to X . Thus, $b_{11} - b_{13} < b_{31} - b_{33}$. ■

Having completed the proof of Theorem 3.1, we now reconsider the theorem in terms of totally positive matrices. A real matrix, finite or infinite, is called *totally positive* (or sometimes *totally nonnegative*) if each $k \times k$ minor is nonnegative. Totally positive matrices have many interesting properties [1, 2] and arise frequently in combinatorics. (See [5, 8, 9, 18, 19, 21].)

One important property of a finite square totally positive matrix is that all of its eigenvalues are nonnegative and real. (See [2, Theorem 1.1]). It is well known that the antiadjacency matrix of any $(\mathbf{2} + \mathbf{2})$ -free, $(\mathbf{3} + \mathbf{1})$ -free poset is totally positive, provided the poset is labelled as in Theorem 1.1. An infinite matrix which is well known to be totally positive is

$$C = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdot & \cdot & \cdot \\ 2 & 1 & 0 & 0 & \cdot & \cdot & \cdot \\ 3 & 2 & 1 & 0 & \cdot & \cdot & \cdot \\ 4 & 3 & 2 & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \end{bmatrix}.$$

It is easy to see that the matrices satisfying the conditions of Theorem 3.1 are essentially submatrices of C : each is determined by a finite multiset of columns and a finite multiset of rows. (See Example 3.1.) Thus, the squared antiadjacency matrix $B = A^2$ of any $(\mathbf{3} + \mathbf{1})$ -free poset is totally positive.

4. CHAIN POLYNOMIALS AND OPEN QUESTIONS

Many open problems in algebraic combinatorics concern the f -vectors of posets and simplicial complexes. (See, for example, [16, Chap. 2–3].) We define the f -vector of a finite poset P to be the integer sequence

$$f_P = (f_0, \dots, f_{d-1}),$$

where f_{i-1} is the number of i -element chains in P , and d is the maximum cardinality of a chain in P . While it would be desirable to characterize the f -vectors corresponding to particular classes of posets, few results of this type are known. Typical results relate the integers f_0, \dots, f_{d-1} by linear and quadratic inequalities. (See [3].) The f -vector is called *unimodal* if

$$f_0 \leq \dots \leq f_j \geq \dots \geq f_{d-1},$$

for some index j , and *log concave* if

$$f_i^2 \geq f_{i-1} f_{i+1},$$

for all $i = 1, \dots, d-2$. To prove that f_P is unimodal and log concave, it suffices to show that the related *chain polynomial*,

$$f_P(x) = 1 + \sum_{i=1}^d f_{i-1} x^i,$$

has only real zeros. Further, the following identity relates the chain polynomial $f_P(x)$ to the antiadjacency matrix A [15],

$$f_P(x) = \det(I + xA). \quad (4.1)$$

Thus, $f_P(x)$ has only real zeros if and only if A has only real eigenvalues.

By Theorem 1.1, the chain polynomial of a $(2+2)$ -free, $(3+1)$ -free poset has only real zeros. (See discussion following Proposition 3.5.) Similarly, by Theorem 3.1, the same holds for $(3+1)$ -free posets. This result was originally shown in [17, Corollary 2.9], using facts about symmetric functions [10, 17, Theorem 2.8].

COROLLARY 4.1. *Let P be a $(3+1)$ -free poset. Then the chain polynomial $f_P(x)$ has only real zeros.*

Proof. Label P as in Theorem 3.1, and let A be the corresponding antiadjacency matrix. By the discussion following Proposition 3.5, the matrix $B = A^2$ is totally positive and therefore has only nonnegative real eigenvalues. It follows that A has only real eigenvalues, and that $f_P(x) = \det(I + xA)$ has only real zeros. ■

The converse of Corollary 4.1 is not true, for there are many posets containing $3+1$ as an induced subposet, whose chain polynomials have only real zeros. An important open problem is to determine which posets have this property. In particular, we have the following conjecture due to Stanley [18] and Neggers [11].

Conjecture 4.2. *Let $J(Q)$ be a finite distributive lattice. Then the chain polynomial $f_{J(Q)}(x)$ has only real zeros.*

Various proofs show that the conjecture holds for the special cases in which Q is a disjoint sum of chains [12], a Ferrers poset [4], and a series-parallel poset [22]. In addition, Stembridge has verified the conjecture for all posets Q having eight or fewer elements [20]. A more general open problem is to determine whether the conjecture holds for the larger class of

modular lattices. No counterexamples are known. It would be interesting to apply the identity (4.1) to either open question or to the special cases.

Another open problem is to count the $(\mathbf{3} + \mathbf{1})$ -free posets of cardinality n . By the comment following Theorem 1.1, the Catalan numbers count $(\mathbf{2} + \mathbf{2})$ -free, $(\mathbf{3} + \mathbf{1})$ -free posets. It would be interesting to apply the discussion following Proposition 3.5 to obtain a simple formula for $(\mathbf{3} + \mathbf{1})$ -free posets.

ACKNOWLEDGMENTS

I thank Richard Stanley for inspiration and for many helpful suggestions. I also thank my referees for suggestions which have simplified several proofs.

REFERENCES

1. M. Aissen, I. J. Schoenberg, and A. Whitney, On generating functions of totally positive sequences, *J. Anal. Math.* **2** (1952), 93–103.
2. T. Ando, Totally positive matrices, *Linear Algebra Appl.* **90** (1987), 165–219.
3. L. J. Billera and G. Hetyei, Linear inequalities for flags in graded partially ordered sets, to appear.
4. F. Brenti, Unimodal, log-concave, and Pólya frequency sequences in combinatorics, *Mem. Amer. Math. Soc.* **413** (1989).
5. F. Brenti, Combinatorics and total positivity, *J. Combin. Theory Ser. A* **71** (1995), 175–218.
6. P. C. Fishburn, Intransitive indifference with unequal indifference intervals, *J. Math. Psych.* **7** (1970), 144–149.
7. P. C. Fishburn, “Interval Graphs and Interval Orders,” Wiley, New York, 1985.
8. S. Fomin and A. Zelevinsky, Double bruhat cells and total positivity, *J. Amer. Math. Soc.* **12** (1999), 335–380.
9. S. Fomin and A. Zelevinsky, Total positivity: Tests and parametrizations, *Math. Intelligencer*, in press.
10. V. Gasharov, Incomparability graphs of $(\mathbf{3} + \mathbf{1})$ -free posets are s -positive, *Discrete Math.* **157** (1996), 211–215.
11. J. Neggers, Representations of finite partially ordered sets, *J. Combin. Inform. System Sci.* **3** (1978), 113–133.
12. R. Simion, A multiindexed Sturm sequence of polynomials and unimodality of certain combinatorial sequences, *J. Combin. Theory Ser. A* **36** (1984), 15–22.
13. R. Stanley, “Enumerative Combinatorics,” Vol. 1, Wadsworth & Brooks/Cole, Belmont, CA, 1986.
14. R. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, *Adv. Math.* **111** (1995), 166–194.
15. R. Stanley, A matrix for counting paths in acyclic digraphs, *J. Combin. Theory Ser. A* **74** (1996), 169–172.
16. R. Stanley, “Combinatorics and Commutative Algebra,” Birkhäuser, Boston, MA, 1996.
17. R. Stanley, Graph colorings and related symmetric functions: Ideas and applications, *Discrete Math.* **193** (1998), 267–286.

18. R. Stanley, Positivity problems and conjectures, in “Mathematics: Frontiers and Perspectives,” International Mathematics Union, in press.
19. R. Stanley and J. R. Stembridge, On immanants of Jacobi–Trudi matrices and permutations with restricted positions, *J. Combin. Theory Ser. A* **62** (1993), 261–279.
20. J. Stembridge, personal communication.
21. J. Stembridge, Immanants of totally positive matrices are nonnegative, *Bull. London Math. Soc.* **23** (1991), 422–428.
22. D. Wagner, Enumeration of functions from posets to chains, *European J. Combin.* **13** (1992), 313–324.